

Welcome to STN International! Enter x:x

LOGINID:sssptal656xw3

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Apr 03 "Ask CAS" for self-help around the clock
NEWS 3 Apr 09 REILSTEIN: Reload and Implementation of a New Subject Area
NEWS 4 Apr 09 ZDB will be removed from STN
NEWS 5 Apr 10 US Patent Applications available in IFICDB, IFIPAT, and
IFIUDB
NEWS 6 Apr 10 Records from IP.com available in CAPLUS, HCAPLUS, and
ZCAPLUS
NEWS 7 Apr 10 BIOSIS Gene Names now available in TOXCENTER
NEWS 8 Apr 10 Federal Research in Progress (FERIP) now available
NEWS 9 Jun 03 New e-mail delivery for search results now available
NEWS 10 Jun 10 MEDLINE Reload
NEWS 11 Jun 10 NCTFULL has been reloaded
NEWS 12 Jul 01 HOPEGE no longer contains STANDARDS file segment
NEWS 13 Jul 27 USAN to be reloaded July 28, 2002;
saved answer sets no longer valid
NEWS 14 Jul 29 Enhanced polymer searching in REGISTRY
NEWS 15 Jul 31 NETFIRST to be removed from STN
NEWS 16 Aug 01 CANCERLIT reload
NEWS 17 Aug 01 PHARMAMarketLetter(PHARMAML) - new on STN
NEWS 18 Aug 01 N113 has been reloaded and enhanced
NEWS 19 Aug 10 Aquatic Toxicity Information Retrieval (AQUIRE)
now available on STN
NEWS 20 Aug 14 IFIPAT, IFICDB, and IFIUDB have been reloaded
NEWS 21 Aug 14 The MEDLINE file segment of TOXCENTER has been reloaded
NEWS 22 Aug 20 Sequence searching in REGISTRY enhanced
NEWS 23 Sep 01 CASIO has been reloaded and enhanced
NEWS 24 Sep 10 Experimental properties added to the REGISTRY file
NEWS 25 Sep 10 Indexing added to some pre-1967 records in CA/CAPLUS
NEWS 26 Sep 10 CA Section Thesaurus available in CAPLUS and CA
NEWS 27 Oct 01 CASREACT Enriched with Reactions from 1907 to 1985
NEWS 28 Oct 21 EVENTLINE has been reloaded

NEWS EXPRESS October 14 CURRENT WINDOWS VERSION IS V6.01,
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),
AND CURRENT DISCOVER FILE IS DATED 01 OCTOBER 2002

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (General Information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer Agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may

result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 12:12:31 ON 24 OCT 2002

=> FIL BIOSIS MEDLINE EMBASE LIFESCI CAPLUS

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'BIOSIS' ENTERED AT 12:12:43 ON 24 OCT 2002

COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(B)

FILE 'MEDLINE' ENTERED AT 12:12:43 ON 24 OCT 2002

FILE 'EMBASE' ENTERED AT 12:12:43 ON 24 OCT 2002

COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'LIFESCI' ENTERED AT 12:12:43 ON 24 OCT 2002

COPYRIGHT (C) 2002 Cambridge Scientific Abstracts (CSA)

FILE 'CAPLUS' ENTERED AT 12:12:43 ON 24 OCT 2002

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s genetic(w) material or DNA or RNA (2a) extract? or isolat?

2 FILES SEARCHED...

4 FILES SEARCHED...

L1 5813969 GENETIC(W) MATERIAL OR DNA OR RNA (2A) EXTRACT? OR ISOLAT?

=> s l1 (s) column or immobiliz?

L2 805809 L1 (S) COLUMN OR IMMOBILIZ?

=> s l2 (s) label?

L2 18648 L2 (S) LABEL?

=> s l3 (s) radical adj1 mediat?

L4 0 L3 (S) RADICAL ADJ1 MEDIAT?

=> s l3 (s) radical(w) mediat?

L4 4 L3 (S) RADICAL(W) MEDIAT?

=> d 1: 1-4

L1 ANSWER 1 OF 4 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

AN 1999:119009 BIOSIS

DN PREV199900119009

TI Increased DNA oxidation and decreased levels of repair products in Alzheimer's disease ventricular CSF.

AF Lovell, Mark A.; Gabbita, S. Prasad; Markesbery, William R. (1)

JA (1) 101 Sanders-Brown Build., Univ. Kentucky, Lexington, KY 40536-0181

TA

SO Journal of Neurochemistry, (Feb., 1999) Vol. 72, No. 2, pp. 771-776.

ISSN: 0022-3042.

BT Article

LA English
 L5 ANSWER 2 OF 4 MEDLINE
 AN 1999127942 MEDLINE
 DN 99127942 PubMed ID: 9939752
 TI Increased DNA oxidation and decreased levels of repair products in
 Alzheimer's disease ventricular CSF.
 AU Lovell M A; Gabbita S P; Markesbery W R
 CS Sanders-Brown Center on Aging, and Department of Chemistry, University of
 Kentucky, Lexington 40536-0230, USA.
 NC 1901-AG15118 (NIA)
 SP 1901-AG15144 (NIA)
 SO JOURNAL OF NEUROCHEMISTRY, (1999 Feb) 72 (2) 771-6.
 Journal code: 2936190R. ISSN: 0022-3042.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 122902
 ED Entered STN: 19991223
 Last Updated on STN: 19991223
 Entered Medline: 19991211

 L5 ANSWER 3 OF 4 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.
 AN 1999042104 EMBASE
 TI Increased DNA oxidation and decreased levels of repair products in
 Alzheimer's disease ventricular CSF.
 AU Lovell M.A.; Gabbita S.P.; Markesbery W.R.
 CS Dr. W.R. Markesbery, 101 Sanders-Brown Building, University of Kentucky,
 Lexington, KY 40536-0230, United States
 SO Journal of Neurochemistry, 1999) 72/2 (771-776).
 Refs: 44
 ISSN: 0022-3042 CODEN: JONHA
 CY United States
 DT Journal; Article
 FS 003 General Pathology and Pathological Anatomy
 003 Neurology and Neurosurgery
 LA English
 SL English

 L5 ANSWER 4 OF 4 LIFESCI COPYRIGHT 2002 CSA
 AN 1999091917 LIFESCI
 TI Increased DNA Oxidation and Decreased Levels of Repair Products in
 Alzheimer's Disease Ventricular CSF
 AU Lovell, M.A.; Gabbita, S.P.; Markesbery, W.R.*
 CS 101 Sanders-Brown Building, University of Kentucky, Lexington, KY
 40536-0230, USA
 SO Journal of Neurochemistry [J. Neurochem.], (19990200) vol. 72, no. 2, pp.
 771-776.
 ISSN: 0022-3042.
 DT Journal
 FS N3
 LA English
 SL English

AS

---Logging off of STN---

=>
Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
FULL ESTIMATED COST	ENTRY	SESSION
	36.21	36.42

STN INTERNATIONAL LOGOFF AT 12:19:22 ON 24 OCT 2002

AN 1999:526234 CAPLUS

DN 131:308477

TI Fluorescent labelling of closely-spaced aldehydes induced in
DNA by bleomycin-Fe(III)

AU Chakrabarti, S.; Mahmood, A.; Makrigiorgos, G. M.

CS Joint Center for Radiation Therapy and Dana Farber Cancer Institute,
Harvard Medical School, Boston, MA, 02215, USASO International Journal of Radiation Biology (1999), 75(8), 1055-1065
CODEN: IJRBEE; ISSN: 1955-3002

PB Taylor & Francis Ltd.

DT Journal

LA English

AB The purpose of this study was to test the ability of two novel fluorescent reagents fluorescent aldehyde-reactive probe (FARP) and FARPhe, to label aldehyde-contg. sites (principally apurinic sites) generated in DNA by the radiomimetic drug bleomycin, and to use fluorescent energy transfer from FARPhe (donor) to FARP (acceptor) to quantitate such closely-spaced sites. FARPhe, 7-hydroxycoumarin-3-carboxylic acid (((((amino-oxymethyl) carbonyl) hydrazino) carbonyl) ethyl) amide) was synthesized with a protocol similar to the one recently reported for FARP (a fluorescein-based probe). Both FARPhe and FARP form stable covalent bonds with the open-chain aldehydes generated upon acidic depurination of DNA. Plasmid DNA exposed to bleomycin-Fe(III)-ascorbate undergoes extensive strand breakage, and upon subsequent reaction with FARPhe and/or FARP it becomes fluorescently labeled, indicating the generation of aldehyde-contg. sites. The binding of the probes to calf thymus or plasmid DNA results in significant fluorescent energy transfer among closely-spaced fluorophores, as revealed by the fluorescence increase following digestion of fluorescently labeled samples with nuclease P1. The fluorescence quenching is most evident when both FARPhe and FARP are used simultaneously to trap aldehyde sites. When single-stranded oligonucleotides engineered to contain either one or two closely spaced bleomycin binding sites are exposed to bleomycin and then fluorescently labeled, the oligonucleotides demonstrate significantly increased fluorescent energy transfer with two binding sites indicating a dependence of aldehyde site generation and clustering on the local sequence of a single strand. In conclusion, a new detection method for DNA damage induced by bleomycin following fluorescent labeling of aldehyde group-contg. sites (FLAGS) and their clustering via fluorescent energy transfer is demonstrated. The method is applicable to any form of DNA. This work may lead to a general approach for the quantification of multiply damaged sites in DNA, a subset of DNA lesions that may have major biol. significance.

RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>

L3 ANSWER 8 OF 16 MEDLINE DUPLICATE 3
 AN 97105898 MEDLINE
 DN 97105898 PubMed ID: 8948646
 TI Chemical methods of DNA and RNA fluorescent labeling.
 AU Froudnikov D; Mirzakekov A
 CS Engelhardt Institute of Molecular Biology, Moscow, Russia.
 SO NUCLEIC ACIDS RESEARCH, (1996 Nov 15) 24 (22) 4535-42.
 Journal code: CBL; 0411011. ISSN: 0305-1048.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199701
 ED Entered STN: 19970219
 Last Updated on STN: 19970219
 Entered Medline: 19970117
 AB Several procedures have been described for fluorescent labeling of DNA and RNA. They are based on the introduction of aldehyde groups by partial depurination of DNA or oxidation of the 3'-terminal ribonucleoside in RNA by sodium periodate. Fluorescent labels with an **attached** hydrazine group are efficiently coupled with the aldehyde groups and the hydrazone bonds are stabilized by reduction with sodium cyanoborohydride. Alternatively, DNA can be quantitatively split at the depurinated sites with ethylenediamine. The aldimine bond between the aldehyde group in depurinated DNA or oxidized RNA and ethylenediamine is stabilized by reduction with sodium cyanoborohydride and the primary amine group introduced at these sites is used for **attachment** of isothiocyanate or succinimide derivatives of fluorescent dyes. The fluorescent DNA labeling can be carried out either in solution or on a reverse phase **column**. These procedures provide simple, inexpensive methods of multiple **DNA labeling** and of introducing one fluorescent dye molecule per RNA, as well as quantitative DNA fragmentation and incorporation of one label per fragment. These methods of fluorophore **attachment** were shown to be efficient for use in the hybridization of labeled RNA, DNA and DNA fragments with oligonucleotide microchips.

L3 ANSWER 10 OF 16 MEDLINE DUPLICATE 4
 AN 94057384 MEDLINE
 DN 94057384 PubMed ID: 8238885
 TI Biotinylation of **DNA** on membrane **supports**: a procedure
 for preparation and easy control of **labeling** of nonradioactive
 single-stranded **nucleic** acid probes.
 AU Eidenko V V
 CS Department of Immunology, Institute of Transplantology and Artificial
 Organs, Moscow, Russia.
 SO ANALYTICAL BIOCHEMISTRY, (1993 Aug 15) 213 (1) 75-8.
 Journal code: 4NK; 0370535. ISSN: 0003-2697.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199312
 ED Entered STN: 19940117
 Last Updated on STN: 19940117
 Entered Medline: 19931203
 AB We have used M13 single-stranded DNA **bound** by uv to small pieces
 of nylon membrane for the synthesis of biotinylated single-stranded DNA
 probes. The labeling method requires a large fragment of DNA polymerase I
 and random hexanucleotides. There is no need for previous linearization of
 the template. The clean probe is removed from the membrane by a single
 wash step. The synthesized probe is completely free of unincorporated
 precursors. This makes possible the easy control of the reaction of
 incorporation of biotinylated analogues into the probe by simple staining
 on the filter, thus allowing evaluation of the efficiency of labeling. The
 DNA membrane can be stored for reuse. With the procedure described it is
 possible to biotinylate many DNA fragments in parallel, simultaneously
 controlling the efficiency of labeling in a time- and cost-saving manner.